

Nonlinear interaction between electromagnetic fields at high temperature

F. T. Brandt, J. Frenkel

Instituto de Física, Universidade de São Paulo,

São Paulo, 01498 SP, Brasil

J. C. Taylor

Department of Applied Mathematics and Theoretical Physics, University of Cambridge,

Cambridge, CB3 9EW, UK

The electron-positron ‘box’ diagram produces an effective action which is fourth order in the electromagnetic field. We examine the behaviour of this effective action at high-temperature (in analytically continued imaginary-time thermal perturbation theory). We argue that there is a finite, nonzero limit as $T \rightarrow \infty$ (where T is the temperature). We calculate this limit in the nonrelativistic static case, and in the long-wavelength limit. We also briefly discuss the self-energy in 2-dimensional QED, which is similar in some respects.

11.10.W

I. INTRODUCTION

The effective action due to the electron-positron ‘box’ diagram was calculated in 1950 [1]. It is fourth order in the electromagnetic field. The object of this paper is to investigate the behaviour of the QED effective action at temperatures high compared to the electron mass and to the frequencies and wave numbers of the electromagnetic field (see also [2]). We use

analytically continued imaginary-time thermal perturbation theory (which is known [3,4] to be related to real-time theory). The question which we seek to answer is: what is the behaviour at high-temperature? Individual contributions contain T^2 and $\log(T)$ terms, where T is the temperature (T^3 and T terms cancel by symmetry). However, gauge invariance imposes strong constraints (just as it does at zero temperature, forbidding UV divergence for example), and one expects strong cancellations. In fact we are able to confirm explicitly the cancellation of the T^2 terms. Note that the dependance upon T for high T is not necessarily simply related to the UV divergence or convergence of the zero-temperature amplitude. In QCD, for example, the n -gluon functions, for all n , go as T^2 although they are UV finite for $N > 4$.

The main concern of this paper is possible $\log(T)$ terms. We have not managed to demonstrate their absence by explicit computation. Instead we give an indirect argument. This (Sec. IV) is based upon first studying the static case and then using Lorentz invariance.

We therefore expect the effective action to have a finite, non-zero limit as $T \rightarrow \infty$. Like the zero-temperature action, this must be gauge-invariant; but it may be more complicated because it is not necessarily Lorentz-invariant. We have not been able to compute it in general. In two special cases, the nonrelativistic static case (which is just a number) and the long wavelength limit (see the Appendix), we have been able to compute the high temperature limit. We have also checked by direct calculation the cancellation of $\log(T)$ terms in the special case in which all the wave-vectors are parallel.

There is an example which has some points of resemblance. That is the self-energy function in 2-dimensional QED. Here (although the zero-temperature function is UV finite) it is not immediately obvious whether there is a $\log(T)$ term. We find, in fact, that there is a finite limit at high T .

Four-photons graphs with two thermal photons and non-thermal fermions have been considered as subdiagrams contributing in higher order to the photon self energy [5]. These lead in the low temperature limit, to temperature-dependent corrections to the dielectric constant and speed of light behaving like $(T/m)^4$. As a possible application of our four-

photon result, one could thermalise two photons obtaining, following [5], a higher order correction to the photon self-energy at high temperature.

In thermal QCD, it is known that a study of the high-temperature behaviour of thermal graphs is an important tool in resumming perturbation theory. In QED, power dependence on T is confined to self-energy graphs. The next question to ask is whether there are graphs with a $\log(T)$ behaviour at large T . This paper addresses that problem.

II. THE THERMAL PERTURBATION THEORY

The graphs which contribute to the nonlinear electromagnetic interaction are shown in Fig. 1. There are three other contributions obtained by charge conjugation.




FIG. 1. Box diagrams which contribute to the nonlinear electromagnetic interaction. Dotted lines represent photons, and solid lines stand for electrons.

The analytically continued imaginary-time thermal perturbation theory can be formulated [6,7] so as to express the thermal amplitude (having subtracted the zero-temperature part) in terms of amplitudes for forward scattering of an electron in an external electromagnetic field, as depicted in Fig. 2. There are 24 Feynman diagrams like this one, which are obtained by even and odd permutations of the external momenta and polarizations. The corresponding analytic expression has the form

$$\frac{e^4}{(2\pi)^3} \int_0^\infty \frac{q^2}{2Q^0} dq N(Q^0) \int d\Omega \sum_{ijkl} B_{(ijkl)}^{\mu\nu\lambda\sigma}(k_1, k_2, k_3; Q). \quad (2.1)$$

Here $q = |\mathbf{Q}|$, $Q^0 = (q^2 + m^2)^{1/2}$, $N(Q^0) = (1 + e^{Q^0/T})^{-1}$, $\int d\Omega$ is an integral over the directions of $\mathbf{Q}$, and the sum is over the permutation $(ijkl)$ of (1234) . Each B has a numerator which is a Dirac trace containing a projection operator $(\mathcal{Q} + m)$. For example

$$B_{(1234)}^{\mu\nu\lambda\sigma} = \frac{\text{tr} \left[(\mathcal{Q} + m) \gamma^\mu (\mathcal{Q} + k_1 + m) \gamma^\nu (\mathcal{Q} + k_{12} + m) \gamma^\lambda (\mathcal{Q} + k_{123} + m) \gamma^\sigma \right]}{(2Q \cdot k_1 + k_1^2) (2Q \cdot k_{12} + k_{12}^2) (2Q \cdot k_{123} + k_{123}^2)}, \quad (2.2)$$

where $k_{12} = k_1 + k_2$, etc. In Eq. 2.1, the k_i^0 are to be given (small) imaginary parts. Different choices of signs for these imaginary parts define the different branches of the analytically continued imaginary-time amplitude [3]. The explicit result for the trace in Eq. (2.2) is given in Eq. (A1) of the Appendix.

We may alternatively write the sum in Eq. (2.1) as

$$\sum_{(ijkl)'} \left[B_{(ijkl)}^{\mu\nu\lambda\sigma}(k_1, k_2, k_3; Q) + B_{(ijkl)}^{\mu\nu\lambda\sigma}(k_1, k_2, k_3; -Q) \right], \quad (2.3)$$

where the $(ijkl)'$ in the summation indicates that only even permutations of (1234) are to be included.




FIG. 2. One of the four diagrams corresponding to the first diagram in Fig. 1. Dotted lines represent photons, and solid lines stand for electrons.

III. THE HIGH-TEMPERATURE BEHAVIOUR

This section is included for completeness. Its conclusion is in fact contained, as the abelian special case, in reference [8]. The straightforward way to get the high-temperature limit of Eq. (2.1) is to use the expansion

$$(2Q \cdot k + k^2)^{-1} = (2Q \cdot k)^{-1} - k^2 (2Q \cdot k)^{-2} + k^4 (2Q \cdot k)^{-3} + \dots \quad (3.1)$$

in the denominators in Eq. (2.2), and to expand the numerator similarly in terms of the k_i . One thus gets in Eq. (2.2) terms which are homogeneous in Q of degrees 1, 0, -1 , -2 . (The expansion (3.1) cannot be taken further without introducing specious infra-red divergences.) The terms of odd degree cancel in Eq. (2.3).

The terms of degree 0 would produce terms $\mathcal{O}(T^2)$ in (1). However, such terms cancel in the sum in Eq. (2.3). As an example, take the terms proportional to $Q^\mu Q^\nu Q^\lambda Q^\sigma$. We find the following contributions (writing $K_i = Q \cdot k_i$, etc):

$$\begin{aligned} & \frac{k_1^2}{K_1^2} \left[\frac{1}{K_{23}} \left(\frac{1}{K_2} + \frac{1}{K_3} \right) + \frac{1}{K_{34}} \left(\frac{1}{K_3} + \frac{1}{K_4} \right) + \frac{1}{K_{42}} \left(\frac{1}{K_4} + \frac{1}{K_2} \right) \right] + (3 \text{ cyclic perms.}) \\ & - \frac{k_{12}^2}{K_{12}^2} \left(\frac{1}{K_3} + \frac{1}{K_4} \right) \left(\frac{1}{K_1} + \frac{1}{K_2} \right) + (2 \text{ cyclic perms. of } 2,3,4) \\ & = \frac{1}{K_1 K_2 K_3 K_4} (k_1^2 + k_2^2 + k_3^2 + k_4^2 - k_{12}^2 - k_{13}^2 - k_{14}^2) = 0. \end{aligned} \quad (3.2)$$

For the terms of degree -2 , there is no such cancellation at the integrand level in general. So, to examine possible $\log(T)$ terms, we have had to resort to an indirect argument. This we present in the next two sections.

IV. ABSENCE OF LOGARITHMIC TEMPERATURE DEPENDENCE

We begin the argument by considering the static case, $k^{0i} = 0$, $i = 1, 2, 3, 4$. This is simpler than the general case, because no analytic continuation is necessary, and so we can write an explicit general formula for the thermal box diagrams. This has the form (unlike in Eq. (2.1), we now refer to the complete thermal amplitude, including the zero-temperature piece)

$$T \sum_{Q^0 = \pi i T (2n+1)} \frac{1}{(2\pi)^3} \int d^{3-\epsilon} \mathbf{Q} F^{\mu\nu\lambda\sigma}(\mathbf{k}_i, \mathbf{Q}, Q^0), \quad (4.1)$$

where n runs over the integers. For fixed n , the $\mathbf{Q}$ -integral is IR and UV finite (i.e. has no pole at $\epsilon = 0$). For

$$T \gg |\mathbf{k}_i|, m, \quad (4.2)$$

Eq. (4.1) gives

$$\sum_n \left[a(\epsilon) T^{-\epsilon} (2n+1)^{-1-\epsilon} + O(T^{-1-\epsilon} n^{-2-\epsilon}) \right]. \quad (4.3)$$

If $a(0)$ is nonzero, this sum diverges; and the corresponding zero-temperature Euclidean field theory (with $\int_{-\infty}^{i\infty} dQ^0$ instead of the sum) would be UV divergent, which we know is not the case. It follows that

$$a(\epsilon) \sim \epsilon b(\mathbf{k}_i/m). \quad (4.4)$$

We may evaluate the sum in (4.1), then let $\epsilon \rightarrow 0$ and finally let $T \rightarrow \infty$ to get a finite result.

As an example, let us consider the special case when $|\mathbf{k}^i| \ll m \ll T$. Then we may write (4.1) in the form

$$\int \frac{d^3 \mathbf{Q}}{(2\pi)^3} \int \frac{dz}{2\pi i} N(z) \frac{N^{\mu\nu\lambda\sigma}(\mathbf{k}_i = 0, \mathbf{Q}, z)}{(z^2 - q^2)^4}, \quad (4.5)$$

where $N^{\mu\nu\lambda\sigma}$ is the numerator of $F^{\mu\nu\lambda\sigma}$. First take the case $\mu = \nu = \lambda = \sigma = 0$. Then all 3 graphs in Fig. 1, together with the ones obtained by charge conjugation, contribute the same, and

$$\begin{aligned} N^{0000} &= \text{tr} [(z + \mathbf{Q} \cdot \boldsymbol{\alpha})^4] \\ &= 2 [(z + q)^4 + (z - q)^4], \end{aligned} \quad (4.6)$$

where $\alpha^i \equiv \gamma^i \gamma^0$. So (4.5) gives

$$\begin{aligned} &\frac{6}{\pi^2} \int_0^\infty dq \, q^2 \int \frac{dz}{2\pi i} N(z) \left[\frac{1}{(z - q)^4} + \frac{1}{(z + q)^4} \right] \\ &= \frac{1}{\pi^2} \int_0^\infty dq q^2 [N'''(q) + N'''(-q)] \\ &= -\frac{4}{\pi^2} N(0) = -\frac{2}{\pi^2}. \end{aligned} \quad (4.7)$$

In this special case, a $\log(T)$ term would have shown itself as

$$\int_0^\infty \frac{dq}{q} N(q). \quad (4.8)$$

The IR divergence at the lower limit would have been a consequence of using the expansion (3.1). But the presence of a term like (4.8) would have correctly signalled the presence of a $\log(T)$ term. As it is we do not find a term like (4.8) in (4.5).

The other components, F^{00ij} etc, must vanish in the limit $\mathbf{k}_i \rightarrow 0$ because of gauge invariance.

We now go from the static case to the general case, so far as $\log(T)$ terms are concerned. Terms of degree -2 in Q in the expansion of (2.2) using (3.1) would produce an integral containing

$$\int_0^\infty \frac{dq}{Q^0} N(Q^0) \quad (4.9)$$

(like (4.8)), and thus a $\log(T)$ term. Setting $m = 0$ in the angular integral in Eq. (2.1), because mass terms do not affect the $\log(T)$ contributions, we then obtain an integrand which is homogeneous of degree -2 in the vector $\hat{Q} = (1, \hat{\mathbf{Q}})$. Such integrals have been studied in [8], where it was shown that they could be expressed in the form

$$\sum_a R_a(k_i) M_a(k_i), \quad (4.10)$$

where the R_a are rational, Lorentz covariant tensors and the M_a are a set of four Lorentz invariant functions, with

$$M_1 = \Delta^{-\frac{1}{2}} \arctan \left[\frac{\Delta^{\frac{1}{2}}}{k_2 \cdot k_3} \right], \quad (4.11)$$

where

$$\Delta = k_2^2 k_3^2 - (k_2 \cdot k_3)^2, \quad (4.12)$$

and M_2, M_3 are defined similarly by cyclic permutation of $(1, 2, 3)$. Finally $M_4 = 1$. The important point about (4.10) is its Lorentz covariance, in spite of the fact that the angular integral in (2.1) is not generally a Lorentz invariant process (except when the integrand is homogeneous in $\hat{Q}$ of degree -2).

Because of the importance for our proof of this result about Lorentz invariance, we here insert a simple argument for this. Consider the part of the angular integral in Eq. (2.1) contributing to $\log(T)$ terms, which has the form

$$\int d\Omega B(k_i, \hat{Q}). \quad (4.13)$$

Because of homogeneity property, we can write this as

$$\int d\Omega |\mathbf{Q}|^2 [B(k_i, Q)]_{Q^0=|\mathbf{Q}|}, \quad (4.14)$$

for any value of $|\mathbf{Q}|$. In particular, we may fix $k.Q = 1$ where k is an arbitrary combination of the k_i . Thus we get

$$2 \int d^4Q \delta(Q^2) \theta(Q^0) \delta(k.Q - 1) B(k_i, Q). \quad (4.15)$$

This final form shows that the integral is a covariant function of the k_i . But the original form shows that it also independent of the particular choice of k . Thus the integral is covariant, as claimed.

We now return to the form (4.10). Clearly the functions defined in (4.11) have branch cuts at $k_i^2 = 0$, and the coefficient rational functions R_a in (4.10) have poles where subsets of the k_i are linearly dependent. In order to avoid these complications, we now assume that all the k_i^0 are pure imaginary, i.e. we are dealing with (4-dimensional) Euclidean field theory. If we can show that there is no $\log(T)$ term in this case, then there can be no such term when we analytically continue further towards the real k_i^0 axes.

Now suppose that there was a $\log(T)$ term in the non-static case; so that not all the R_a in (4.10) were zero. We shall show that this would contradict our previous conclusion that there was no $\log(T)$ term in the static case. Since (4.10) is covariant under rotations in 4-dimensional Euclidean space, we may go to a coordinate system in which the (imaginary) time axis is perpendicular to each of the k_i . This is just the static case ($k_i^0 = 0$). Thus the nonvanishing of (4.10) would contradict what we know about the absence of $\log(T)$ terms in that case. This completes our argument about the general absence of $\log(T)$ terms.

V. DISCUSSION

In order to inquire further the absence of $\log(T)$ terms in the box diagram, we now briefly consider the photon self-energy in two-dimensional QED, which is similar in the

sense that there is no UV divergence at zero-temperature. Of course, the main difference is the role of the axial anomaly in two-dimensional QED [9], which has no counterpart in the four-dimensional box diagram.

Since this model is much simpler, we can give an explicit form for the self-energy at infinite temperature. By using a method similar to that in Sec. IV, the high-temperature limit is found simply by setting

$$N(z) \rightarrow \frac{1}{2} \quad (5.1)$$

in the analogue of Eq. (4.5), evaluated for general momenta k_μ . Then the result for the complete self-energy at infinite temperature is

$$\frac{e^2}{\pi} \left[\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right]. \quad (5.2)$$

Note that the zero-temperature result is [10]

$$\frac{e^2}{\pi} \left[1 + \frac{4m^2}{k^2 R} \arctan(R) \right] \left[\eta^{\mu\nu} - \frac{k^\mu k^\nu}{k^2} \right], \quad (5.3)$$

where $k^2 < 4m^2$ and $R = [4m^2/k^2 - 1]^{1/2}$. Thus the effect of the additional thermal contributions (at high T) is simply to cancel the m^2 term, leaving just the anomaly term behind.

An interesting question concerns the thermal behavior of QED Green functions with $n > 4$ external photon lines (for n -odd, these vanish identically by charge conjugation). By power counting, these functions are UV finite at zero temperature. Hence, using arguments like those in Sec. IV, we see that in the static case the $\log(T)$ terms will be absent. Furthermore, from the work in references [8,11], it follows that the T^2 contributions will cancel out. Therefore, we conclude that in the static case these Green functions will have a finite limit as $T \rightarrow \infty$. Whether this continues to be true in general still remains to be investigated.

ACKNOWLEDGMENTS

F.T.B. and J.F. thank to CNPQ, Brasil, for a grant.

APPENDIX A:

In this appendix we present another explicit result for the high-temperature behavior of the nonlinear electromagnetic interaction. Since in Sec. IV we have shown that there is no $\log(T)$ contribution and computed the static case, it would be also interesting to know the explicit expression for the *long wavelength limit*.

Our starting point is Eq. (2.2). The computation of the trace is a tedious but straightforward task which can be accomplished using a computer program for symbolic manipulations. The result is

$$\begin{aligned}
& 4 (\ k_1^\sigma k_{12}^\lambda k_{123}^\nu Q^\mu - k_1^\lambda k_{12}^\sigma k_{123}^\nu Q^\mu + k_1^\sigma k_{12}^\nu k_{123}^\lambda Q^\mu + k_1^\nu k_{12}^\sigma k_{123}^\lambda Q^\mu + \\
& \ k_1^\lambda k_{12}^\nu k_{123}^\sigma Q^\mu + k_1^\nu k_{12}^\lambda k_{123}^\sigma Q^\mu - k_1^\sigma k_{12}^\lambda k_{123}^\mu Q^\nu + k_1^\lambda k_{12}^\sigma k_{123}^\mu Q^\nu - \\
& \ k_1^\sigma k_{12}^\mu k_{123}^\lambda Q^\nu + k_1^\mu k_{12}^\sigma k_{123}^\lambda Q^\nu - k_1^\lambda k_{12}^\mu k_{123}^\sigma Q^\nu + k_1^\mu k_{12}^\lambda k_{123}^\sigma Q^\nu + \\
& \ 2 k_{12}^\sigma k_{123}^\lambda Q^\mu Q^\nu + 2 k_{12}^\lambda k_{123}^\sigma Q^\mu Q^\nu - k_1^\sigma k_{12}^\nu k_{123}^\mu Q^\lambda - k_1^\nu k_{12}^\sigma k_{123}^\mu Q^\lambda + \\
& \ k_1^\sigma k_{12}^\mu k_{123}^\nu Q^\lambda - k_1^\mu k_{12}^\sigma k_{123}^\nu Q^\lambda + k_1^\nu k_{12}^\mu k_{123}^\sigma Q^\lambda + k_1^\mu k_{12}^\nu k_{123}^\sigma Q^\lambda + \\
& \ 2 k_1^\sigma k_{123}^\nu Q^\mu Q^\lambda - 2 k_{12}^\sigma k_{123}^\nu Q^\mu Q^\lambda + 2 k_1^\nu k_{123}^\sigma Q^\mu Q^\lambda + 2 k_{12}^\nu k_{123}^\sigma Q^\mu Q^\lambda - \\
& \ 2 k_1^\sigma k_{123}^\mu Q^\nu Q^\lambda + 2 k_1^\mu k_{123}^\sigma Q^\nu Q^\lambda + 4 k_{123}^\sigma Q^\mu Q^\nu Q^\lambda + k_1^\lambda k_{12}^\nu k_{123}^\mu Q^\sigma + \\
& \ k_1^\nu k_{12}^\lambda k_{123}^\mu Q^\sigma - k_1^\lambda k_{12}^\mu k_{123}^\nu Q^\sigma + k_1^\mu k_{12}^\lambda k_{123}^\nu Q^\sigma + k_1^\nu k_{12}^\mu k_{123}^\lambda Q^\sigma + \\
& \ k_1^\mu k_{12}^\nu k_{123}^\lambda Q^\sigma + 2 k_1^\lambda k_{12}^\nu Q^\mu Q^\sigma + 2 k_1^\nu k_{12}^\lambda Q^\mu Q^\sigma - 2 k_1^\lambda k_{12}^\nu Q^\mu Q^\sigma + \\
& \ 2 k_{12}^\lambda k_{123}^\nu Q^\mu Q^\sigma + 2 k_1^\nu k_{123}^\lambda Q^\mu Q^\sigma + 2 k_{12}^\nu k_{123}^\lambda Q^\mu Q^\sigma - 2 k_1^\lambda k_{12}^\mu Q^\nu Q^\sigma + \\
& \ 2 k_1^\mu k_{12}^\lambda Q^\nu Q^\sigma + 2 k_1^\lambda k_{123}^\mu Q^\nu Q^\sigma + 2 k_1^\mu k_{123}^\lambda Q^\nu Q^\sigma + 4 k_{12}^\lambda Q^\mu Q^\nu Q^\sigma + \\
& \ 4 k_{123}^\lambda Q^\mu Q^\nu Q^\sigma + 2 k_1^\nu k_{12}^\mu Q^\lambda Q^\sigma + 2 k_1^\mu k_{12}^\nu Q^\lambda Q^\sigma + 4 k_1^\nu Q^\mu Q^\lambda Q^\sigma + \\
& \ 4 k_{12}^\nu Q^\mu Q^\lambda Q^\sigma + 4 k_1^\mu Q^\nu Q^\lambda Q^\sigma + 8 Q^\mu Q^\nu Q^\lambda Q^\sigma + k_1^\sigma k_{12}^\lambda k_{123} \cdot Q \eta^{\mu\nu} - \\
& \ k_1^\lambda k_{12}^\sigma k_{123} \cdot Q \eta^{\mu\nu} + k_1^\sigma k_{12} \cdot Q k_{123}^\lambda \eta^{\mu\nu} - k_1 \cdot Q k_{12}^\sigma k_{123}^\lambda \eta^{\mu\nu} + k_1^\lambda k_{12} \cdot Q k_{123}^\sigma \eta^{\mu\nu} - \\
& \ k_1 \cdot Q k_{12}^\lambda k_{123}^\sigma \eta^{\mu\nu} - k_1^\sigma k_{12} \cdot k_{123} Q^\lambda \eta^{\mu\nu} + k_1 \cdot k_{123} k_{12}^\sigma Q^\lambda \eta^{\mu\nu} - k_1 \cdot k_{123} k_{12}^\sigma Q^\lambda \eta^{\mu\nu} - \\
& \ 2 k_1 \cdot Q k_{123}^\sigma Q^\lambda \eta^{\mu\nu} + k_1^\lambda k_{12} \cdot k_{123} Q^\sigma \eta^{\mu\nu} + 2 k_1^\lambda k_{12} \cdot Q Q^\sigma \eta^{\mu\nu} - k_1 \cdot k_{123} k_{12}^\lambda Q^\sigma \eta^{\mu\nu} - \\
& \ 2 k_1 \cdot Q k_{12}^\lambda Q^\sigma \eta^{\mu\nu} - k_1 \cdot k_{12} k_{123}^\lambda Q^\sigma \eta^{\mu\nu} - 2 k_1 \cdot Q k_{123}^\lambda Q^\sigma \eta^{\mu\nu} - 2 k_1 \cdot k_{12} Q^\lambda Q^\sigma \eta^{\mu\nu} - \\
& \ 4 k_1 \cdot Q Q^\lambda Q^\sigma \eta^{\mu\nu} + k_1^\sigma k_{12}^\nu k_{123} \cdot Q \eta^{\mu\lambda} + k_1^\nu k_{12}^\sigma k_{123} \cdot Q \eta^{\mu\lambda} - k_1^\sigma k_{12} \cdot Q k_{123}^\nu \eta^{\mu\lambda} + \\
& \ k_1 \cdot Q k_{12}^\sigma k_{123}^\nu \eta^{\mu\lambda} - k_1^\nu k_{12} \cdot Q k_{123}^\sigma \eta^{\mu\lambda} - k_1 \cdot Q k_{12}^\nu k_{123}^\sigma \eta^{\mu\lambda} + k_1^\sigma k_{12} \cdot k_{123} Q^\nu \eta^{\mu\lambda} -
\end{aligned}$$

$$\begin{aligned}
& k_1 \cdot k_{123} k_{12}^\sigma Q^\nu \eta^{\mu\lambda} + 2 k_1^\sigma k_{123} \cdot Q Q^\nu \eta^{\mu\lambda} + k_1 \cdot k_{12} k_{123}^\sigma Q^\nu \eta^{\mu\lambda} - k_1^\nu k_{12} \cdot k_{123} Q^\sigma \eta^{\mu\lambda} - \\
& 2 k_1^\nu k_{12} \cdot Q Q^\sigma \eta^{\mu\lambda} - k_1 \cdot k_{123} k_{12}^\nu Q^\sigma \eta^{\mu\lambda} - 2 k_1 \cdot Q k_{12}^\nu Q^\sigma \eta^{\mu\lambda} + k_1 \cdot k_{12} k_{123}^\nu Q^\sigma \eta^{\mu\lambda} + \\
& 2 k_1 \cdot Q k_{123}^\nu Q^\sigma \eta^{\mu\lambda} + 2 k_1 \cdot k_{12} Q^\nu Q^\sigma \eta^{\mu\lambda} - 2 k_1 \cdot k_{123} Q^\nu Q^\sigma \eta^{\mu\lambda} - k_1^\lambda k_{12}^\nu k_{123} \cdot Q \eta^{\mu\sigma} - \\
& k_1^\nu k_{12}^\lambda k_{123} \cdot Q \eta^{\mu\sigma} + k_1^\lambda k_{12} \cdot Q k_{123}^\nu \eta^{\mu\sigma} - k_1 \cdot Q k_{12}^\lambda k_{123}^\nu \eta^{\mu\sigma} - k_1^\nu k_{12} \cdot Q k_{123}^\lambda \eta^{\mu\sigma} - \\
& k_1 \cdot Q k_{12}^\nu k_{123}^\lambda \eta^{\mu\sigma} - k_1^\lambda k_{12} \cdot k_{123} Q^\nu \eta^{\mu\sigma} + k_1 \cdot k_{123} k_{12}^\lambda Q^\nu \eta^{\mu\sigma} - 2 k_1^\lambda k_{123} \cdot Q Q^\nu \eta^{\mu\sigma} + \\
& k_1 \cdot k_{12} k_{123}^\lambda Q^\nu \eta^{\mu\sigma} + k_1^\nu k_{12} \cdot k_{123} Q^\lambda \eta^{\mu\sigma} + k_1 \cdot k_{123} k_{12}^\nu Q^\lambda \eta^{\mu\sigma} - k_1 \cdot k_{12} k_{123}^\nu Q^\lambda \eta^{\mu\sigma} - \\
& 2 k_1 \cdot Q k_{123}^\nu Q^\lambda \eta^{\mu\sigma} + 2 k_1 \cdot k_{123} Q^\nu Q^\lambda \eta^{\mu\sigma} - k_1^\sigma k_{12}^\mu k_{123} \cdot Q \eta^{\nu\lambda} + k_1^\mu k_{12}^\sigma k_{123} \cdot Q \eta^{\nu\lambda} + \\
& k_1^\sigma k_{12} \cdot Q k_{123}^\mu \eta^{\nu\lambda} - k_1 \cdot Q k_{12}^\sigma k_{123}^\mu \eta^{\nu\lambda} - k_1^\mu k_{12} \cdot Q k_{123}^\sigma \eta^{\nu\lambda} + k_1 \cdot Q k_{12}^\mu k_{123}^\sigma \eta^{\nu\lambda} - \\
& k_1^\sigma k_{12} \cdot k_{123} Q^\mu \eta^{\nu\lambda} + k_1 \cdot k_{123} k_{12}^\sigma Q^\mu \eta^{\nu\lambda} - 2 k_1^\sigma k_{123} \cdot Q Q^\mu \eta^{\nu\lambda} + 2 k_{12}^\sigma k_{123} \cdot Q Q^\mu \eta^{\nu\lambda} - \\
& k_1 \cdot k_{12} k_{123}^\sigma Q^\mu \eta^{\nu\lambda} - 2 k_{12} \cdot Q k_{123}^\sigma Q^\mu \eta^{\nu\lambda} - k_1^\mu k_{12} \cdot k_{123} Q^\sigma \eta^{\nu\lambda} - 2 k_1^\mu k_{12} \cdot Q Q^\sigma \eta^{\nu\lambda} + \\
& k_1 \cdot k_{123} k_{12}^\mu Q^\sigma \eta^{\nu\lambda} + 2 k_1 \cdot Q k_{12}^\mu Q^\sigma \eta^{\nu\lambda} - k_1 \cdot k_{12} k_{123}^\mu Q^\sigma \eta^{\nu\lambda} - 2 k_1 \cdot Q k_{123}^\mu Q^\sigma \eta^{\nu\lambda} - \\
& 2 k_1 \cdot k_{12} Q^\mu Q^\sigma \eta^{\nu\lambda} + 2 k_1 \cdot k_{123} Q^\mu Q^\sigma \eta^{\nu\lambda} - 2 k_{12} \cdot k_{123} Q^\mu Q^\sigma \eta^{\nu\lambda} - 4 k_{12} \cdot Q Q^\mu Q^\sigma \eta^{\nu\lambda} + \\
& k_1 \cdot Q k_{12} \cdot k_{123} \eta^{\mu\sigma} \eta^{\nu\lambda} - k_1 \cdot k_{123} k_{12} \cdot Q \eta^{\mu\sigma} \eta^{\nu\lambda} + k_1 \cdot k_{12} k_{123} \cdot Q \eta^{\mu\sigma} \eta^{\nu\lambda} + 2 k_1 \cdot Q k_{123} \cdot Q \eta^{\mu\sigma} \eta^{\nu\lambda} + \\
& k_1^\lambda k_{12}^\mu k_{123} \cdot Q \eta^{\nu\sigma} - k_1^\mu k_{12}^\lambda k_{123} \cdot Q \eta^{\nu\sigma} - k_1^\lambda k_{12} \cdot Q k_{123}^\mu \eta^{\nu\sigma} + k_1 \cdot Q k_{12}^\lambda k_{123}^\mu \eta^{\nu\sigma} - \\
& k_1^\mu k_{12} \cdot Q k_{123}^\lambda \eta^{\nu\sigma} + k_1 \cdot Q k_{12}^\mu k_{123}^\lambda \eta^{\nu\sigma} + k_1^\lambda k_{12} \cdot k_{123} Q^\mu \eta^{\nu\sigma} - k_1 \cdot k_{123} k_{12}^\lambda Q^\mu \eta^{\nu\sigma} + \\
& 2 k_1^\lambda k_{123} \cdot Q Q^\mu \eta^{\nu\sigma} - 2 k_{12}^\lambda k_{123} \cdot Q Q^\mu \eta^{\nu\sigma} - k_1 \cdot k_{12} k_{123}^\lambda Q^\mu \eta^{\nu\sigma} - 2 k_{12} \cdot Q k_{123}^\lambda Q^\mu \eta^{\nu\sigma} + \\
& k_1^\mu k_{12} \cdot k_{123} Q^\lambda \eta^{\nu\sigma} - k_1 \cdot k_{123} k_{12}^\mu Q^\lambda \eta^{\nu\sigma} + k_1 \cdot k_{12} k_{123}^\mu Q^\lambda \eta^{\nu\sigma} + 2 k_1 \cdot Q k_{123}^\mu Q^\lambda \eta^{\nu\sigma} - \\
& 2 k_1 \cdot k_{123} Q^\mu Q^\lambda \eta^{\nu\sigma} + 2 k_{12} \cdot k_{123} Q^\mu Q^\lambda \eta^{\nu\sigma} - k_1 \cdot Q k_{12} \cdot k_{123} \eta^{\mu\lambda} \eta^{\nu\sigma} + k_1 \cdot k_{123} k_{12} \cdot Q \eta^{\mu\lambda} \eta^{\nu\sigma} - \\
& k_1 \cdot k_{12} k_{123} \cdot Q \eta^{\mu\lambda} \eta^{\nu\sigma} - 2 k_1 \cdot Q k_{123} \cdot Q \eta^{\mu\lambda} \eta^{\nu\sigma} - k_1^\nu k_{12}^\mu k_{123} \cdot Q \eta^{\lambda\sigma} - k_1^\mu k_{12}^\nu k_{123} \cdot Q \eta^{\lambda\sigma} + \\
& k_1^\nu k_{12} \cdot Q k_{123}^\mu \eta^{\lambda\sigma} + k_1 \cdot Q k_{12}^\nu k_{123}^\mu \eta^{\lambda\sigma} + k_1^\mu k_{12} \cdot Q k_{123}^\nu \eta^{\lambda\sigma} - k_1 \cdot Q k_{12}^\mu k_{123}^\nu \eta^{\lambda\sigma} - \\
& k_1^\nu k_{12} \cdot k_{123} Q^\mu \eta^{\lambda\sigma} - k_1 \cdot k_{123} k_{12}^\nu Q^\mu \eta^{\lambda\sigma} - 2 k_1^\nu k_{123} \cdot Q Q^\mu \eta^{\lambda\sigma} - 2 k_{12}^\nu k_{123} \cdot Q Q^\mu \eta^{\lambda\sigma} + \\
& k_1 \cdot k_{12} k_{123}^\nu Q^\mu \eta^{\lambda\sigma} + 2 k_{12} \cdot Q k_{123}^\nu Q^\mu \eta^{\lambda\sigma} - k_1^\mu k_{12} \cdot k_{123} Q^\nu \eta^{\lambda\sigma} + k_1 \cdot k_{123} k_{12}^\mu Q^\nu \eta^{\lambda\sigma} - \\
& 2 k_1^\mu k_{123} \cdot Q Q^\nu \eta^{\lambda\sigma} - k_1 \cdot k_{12} k_{123}^\mu Q^\nu \eta^{\lambda\sigma} - 2 k_{12} \cdot k_{123} Q^\mu Q^\nu \eta^{\lambda\sigma} - 4 k_{123} \cdot Q Q^\mu Q^\nu \eta^{\lambda\sigma} + \\
& k_1 \cdot Q k_{12} \cdot k_{123} \eta^{\mu\nu} \eta^{\lambda\sigma} - k_1 \cdot k_{123} k_{12} \cdot Q \eta^{\mu\nu} \eta^{\lambda\sigma} + k_1 \cdot k_{12} k_{123} \cdot Q \eta^{\mu\nu} \eta^{\lambda\sigma} + 2 k_1 \cdot Q k_{123} \cdot Q \eta^{\mu\nu} \eta^{\lambda\sigma})
\end{aligned}
\tag{A1}$$

In the long wavelength limit all wave vectors are set equal to zero and Eq. (A1) simplifies considerably. The denominators in Eq. (2.2) will not depend on $\mathbf{Q}$ and the angular integrals

in (2.1) can be easily computed. Using partial fractions decomposition the result can be expressed in terms of integrals like

$$\int_0^\infty \frac{dq}{\exp(q/T) + 1} \frac{1}{q \pm s}, \quad (\text{A2})$$

where s a linear function of the external frequencies. In the limit $T \rightarrow \infty$ we can replace each individual integral by

$$\int_0^T \frac{dq}{2} \frac{1}{q \pm s} = \frac{1}{2} (\log(T) - \log(\pm s) + \mathcal{O}(1/T)). \quad (\text{A3})$$

When we add together all the contributions, we obtain the result

$$\frac{e^4}{15\pi^2} \left(A(\omega_1, \omega_2, \omega_3, \omega_4) \delta^{ij} \delta^{kl} + A(\omega_1, \omega_3, \omega_2, \omega_4) \delta^{ik} \delta^{jl} + A(\omega_1, \omega_4, \omega_2, \omega_3) \delta^{il} \delta^{jk} \right), \quad (\text{A4})$$

where $\omega_1 + \omega_2 + \omega_3 + \omega_4 = 0$ and the indices (i, j, k, l) are in correspondence with $(\mu, \nu, \lambda, \sigma)$. The function $A(\omega_1, \omega_2, \omega_3, \omega_4)$ is a combination of rational functions of the frequencies times expressions like (A3), and it is symmetric under the permutations $\omega_1 \leftrightarrow \omega_2$, $\omega_3 \leftrightarrow \omega_4$ and $(\omega_1, \omega_2) \leftrightarrow (\omega_3, \omega_4)$.

From (A3) we can see that the function $A(\omega_1, \omega_2, \omega_3, \omega_4)$ grows not faster than $\log(T)$ in the high temperature limit. This is just a special case of the result in Sec. III, where we have shown that the cancellation of contributions proportional to T^2 occurs at the integrand level. We also expect from Sec. IV that the $\log(T)$ contributions should vanish. Indeed, we have obtained that the coefficient of $\log(T)$ is proportional to the following vanishing combination of rational functions

$$\begin{aligned} & c(\omega_1, \omega_2, \omega_3, \omega_4) + c(\omega_2, \omega_1, \omega_3, \omega_4) + c(\omega_3, \omega_4, \omega_1, \omega_2) + c(\omega_4, \omega_3, \omega_1, \omega_2) + \\ & d(\omega_1, \omega_2, \omega_3, \omega_4) + d(\omega_2, \omega_1, \omega_3, \omega_4) + e(\omega_1, \omega_2, \omega_3, \omega_4) = 0 \end{aligned} \quad (\text{A5})$$

where

$$c(\omega_1, \omega_2, \omega_3, \omega_4) = \frac{\omega_1^2 [10 \omega_2^2 \omega_3 \omega_4 + (\omega_1 + 5 \omega_2) (\omega_1 \omega_2 \omega_3 \omega_4 + \omega_1 \omega_3 \omega_4)]}{\omega_{12} \omega_{13} \omega_2 \omega_{23} \omega_3 \omega_4}, \quad (\text{A6a})$$

$$d(\omega_1, \omega_2, \omega_3, \omega_4) = \frac{\omega_{13}^2 (\omega_1^2 + 5 \omega_1 \omega_2 - 3 \omega_1 \omega_3 - 5 \omega_2 \omega_3 - 4 \omega_3^2)}{\omega_1 \omega_2 \omega_3 \omega_4} \quad (\text{A6b})$$

and

$$e(\omega_1, \omega_2, \omega_3, \omega_4) = \frac{(\omega_{12} \omega_{34})^2}{\omega_1 \omega_2 \omega_3 \omega_4}. \quad (\text{A6c})$$

Finally, the function $A(\omega_1, \omega_2, \omega_3, \omega_4)$ is given by

$$\begin{aligned} A(\omega_1, \omega_2, \omega_3, \omega_4) = & c(\omega_1, \omega_2, \omega_3, \omega_4) \log\left(-\frac{\omega_1^2}{\omega_{12}\omega_{34}}\right) + c(\omega_2, \omega_1, \omega_3, \omega_4) \log\left(-\frac{\omega_2^2}{\omega_{12}\omega_{34}}\right) + \\ & c(\omega_3, \omega_4, \omega_1, \omega_2) \log\left(-\frac{\omega_3^2}{\omega_{12}\omega_{34}}\right) + c(\omega_4, \omega_3, \omega_1, \omega_2) \log\left(-\frac{\omega_4^2}{\omega_{12}\omega_{34}}\right) + \\ & d(\omega_1, \omega_2, \omega_3, \omega_4) \log\left(\frac{\omega_{13}\omega_{24}}{\omega_{12}\omega_{34}}\right) + d(\omega_2, \omega_1, \omega_3, \omega_4) \log\left(\frac{\omega_{23}\omega_{14}}{\omega_{12}\omega_{34}}\right). \end{aligned} \quad (\text{A7})$$

The symmetries of $A(\omega_1, \omega_2, \omega_3, \omega_4)$ under permutations $\omega_1 \leftrightarrow \omega_2$, $\omega_3 \leftrightarrow \omega_4$ and $(\omega_1, \omega_2) \leftrightarrow (\omega_3, \omega_4)$, can be easily verified using momentum conservation and Eqs. (A6). One can also notice that $A(\omega_1, \omega_2, \omega_3, \omega_4)$ is real when all the frequencies are real.

- [1] R. Karplus and M. Neuman, Phys Rev. **80**, 380 (1950); **83**, 776 (1951).
- [2] W. Dittrich, Phys. Rev. D **19**, 2385 (1979); M. Loewe and J. C. Rojas, Phys. Rev. D **46**, 2689 (1992).
- [3] T. S. Evans, Nucl. Phys. **B374**, 340 (1992).
- [4] J. C. Taylor, Phys. Rev. D **47**, 725 (1993).
- [5] R. Tarrach, Phys Lett. **133B**, 259 (1983).
- [6] G. Barton, Ann. Phys. **200**, 271 (1990).
- [7] J. Frenkel and J. C. Taylor, Nucl. Phys. **B374**, 156 (1992).
- [8] J. Frenkel and J. C. Taylor, Nucl. Phys. **B334**, 199 (1990).
- [9] R. Baier and E. Pilon, Z. Phys. C52, 339 (1991).
- [10] C. Adam, R.A. Bertlmann and P. Hofer, Riv. Nuovo Cim. 16-no.8, 1 (1993).
- [11] E. Braaten and R. D. Pisarski, Nucl. Phys. **B339**, 310 (1990).